

Complete population transfer in a degenerate 3-level atom

Kh.Yu. Rakhimov*, J.H. McGuire, and Kh.Kh. Shakov

Physics Department, Tulane University, New Orleans, LA 70118 USA.

(Dated: February 1, 2008)

Abstract

We find conditions required to achieve complete population transfer, via coherent population trapping, from an initial state to a designated final state at a designated time in a degenerate 3-level atom, where transitions are caused by an external interaction. Complete population transfer from an initially occupied state 1 to a designated state 2 occurs under two conditions. First, there is a constraint on the ratios of the transition matrix elements of the external interaction. Second, there is a constraint on the action integral over the interaction, or "area", corresponding to the phase shift induced by the external interaction. Both conditions may be expressed in terms of simple odd integers.

PACS numbers: 32.80.-t, 42.50.-p, 32.80.Qk

* Permanent address: Department of Heat Physics, Uzbekistan Academy of Sciences, 28 Katartal St., Tashkent 700135, Uzbekistan

I. INTRODUCTION

Population control in quantum systems, namely transfer of electrons from an ensemble of atoms all in the same initial state to specified final states, is used in problems ranging from coherent population trapping [1, 2, 3, 4, 5], including electromagnetically induced transparency [6, 7, 8, 9], and quantum computing [10, 11, 12, 13, 14, 15, 16], to chemical dynamics [17, 18]. These problems are modeled in terms of an n -level (often 3-level) atom interacting with a strong external field [19, 20, 21, 22, 23, 24]. In this paper we consider population control in a nearly degenerate 3-level atom. We show that the 3-level atom is relatively easy to understand in the degenerate limit, where all 3 states have the same energy. We specifically show how to achieve complete population transfer in a degenerate 3-level atom where the matrix elements of the external interaction have a common time dependence.

Our degenerate approximation [25] is mathematically similar to the rotating wave approximation (RWA) [19, 22], which has been widely applied to both 2 and 3-level atomic models. However, in the RWA degenerate atomic states are not used. Instead one tunes the frequency of the external field to the frequency difference of two non-degenerate levels so that the detuning parameter [20] tends to zero. Thus, in RWA the initial state of an atom plus one photon is degenerate in energy with the final state of the atom. One advantage of using degenerate atomic states, as done in this paper, is that one may use external interactions with a broad range of frequencies. Another advantage is that the interaction frequency can be used as a control parameter, e.g. to vary the duration of time that the transferred population remains in the designated state, or to reduce the population leakage that occurs when the three levels are not fully degenerate.

In the next section we derive analytic formulae for the probabilities, $P_{1,2,3}(t)$, that the electron is in level 1, 2 or 3 at time t . Then we seek conditions for complete population transfer to a designated level that is initially unoccupied at a designated time t_0 . This places conditions on both the relative strengths of the interaction matrix elements, $V_{ij}(t)$, and the action integral, $A(t_0) = \int_0^{t_0} V(t')dt'$. We show that these conditions may be expressed in terms of two odd integers, n_1 and n_2 . Some analysis is done for population leakage, when the levels are not quite degenerate. Calculations are presented.

II. THEORY

Let us consider an n -level atom interacting with an external field, $V_{ext}(\vec{r}, t)$. The total Hamiltonian for this system is $H = H_0 + V_{ext}(\vec{r}, t)$. The n eigenstates, ϕ_k , and corresponding eigenenergies, E_k , of H_0 are assumed to be known. The total wavefunction may be expanded in terms of the known eigenstates, namely, $\Psi(t) = a_1(t)\phi_1 + a_2(t)\phi_2 + \cdots + a_n(t)\phi_n$. With atomic units, using $i\dot{\Psi} = (H_0 + V_{ext}(\vec{r}, t))\Psi$, with $H_0\phi_k = E_k\phi_k$ and $\int \phi_j^*\phi_k d\vec{r} = \delta_{jk}$, one then obtains [20],

$$i\dot{a}_j(t) = E_j a_j(t) + \sum_{k=1}^n V_{jk}(t) a_k(t) \quad , \quad (1)$$

where $V_{jk}(t) = \int \phi_j^* V_{ext}(\vec{r}, t) \phi_k d\vec{r}$. These equations are exact for an n -level atom.

We now require that the system be degenerate, namely that all the energies, E_j , are the same. Since the zero point of energy is arbitrary, one may generally set $E_j = 0$. These conditions give the coupled equations for a degenerate n -level system, namely,

$$i\dot{a}_j(t) = \sum_{k=1}^n V_{jk}(t) a_k(t) \quad . \quad (2)$$

We use the initial condition $a_1(0) = 1$, and $a_j(0) = 0$ for $j \neq 1$. We additionally require that all of the $V_{jk}(t)$ have the same time dependence. Here we also take $V_{jk} = V_{kj}$ to be real.

A. Degenerate 3-level atom

The degenerate 3-level atom can be solved analytically. Recalling that the $V_{jk}(t)$ have a common time dependence, we choose $V(t) = V_{23}(t) = \alpha^{-1}V_{12}(t) = \beta^{-1}V_{13}(t) = \epsilon_j^{-1}V_{jj}(t)$. A relatively simple solution, presented next, is found by taking $\epsilon_j = 0$. More general solutions for $\epsilon_j \neq 0$ are discussed in an appendix. With $\epsilon_j = 0$ and $n = 3$, Eq(2) becomes,

$$\begin{aligned} i\dot{a}_1(t) &= \alpha V(t) a_2(t) + \beta V(t) a_3(t) \quad , \\ i\dot{a}_2(t) &= \alpha V(t) a_1(t) + V(t) a_3(t) \quad , \\ i\dot{a}_3(t) &= \beta V(t) a_1(t) + V(t) a_2(t) \quad . \end{aligned} \quad (3)$$

Now consider the linear combination $c(t) = a_1(t) + x a_2(t) + y a_3(t)$, where x and y are

some time-independent coefficients that we next determine. Then,

$$\begin{aligned}
i\dot{c}(t) &= i\dot{a}_1(t) + x i\dot{a}_2(t) + y i\dot{a}_3(t) \\
&= \alpha V(t) a_2(t) + \beta V(t) a_3(t) + x [\alpha V(t) a_1(t) + V(t) a_3(t)] \\
&\quad + y [\beta V(t) a_1(t) + V(t) a_2(t)] .
\end{aligned} \tag{4}$$

Set $z = \alpha x + \beta y$. Then, $i\dot{c}(t) = zV(t)[a_1(t) + (\frac{\alpha+y}{\alpha x + \beta y})a_2(t) + (\frac{\beta+x}{\alpha x + \beta y})a_3(t)]$. We require that $i\dot{c}(t) = zV(t)c(t)$. This holds if and only if $x = (\alpha + y)/(\alpha x + \beta y)$ and $y = (\beta + x)/(\alpha x + \beta y)$. After some algebra this leads to the useful cubic equation,

$$(\beta^2 - \alpha^2)y^3 + \alpha(2 - \alpha^2 - \beta^2)y^2 + (2\alpha^2 - \beta^2 - 1)y + \alpha(\beta^2 - 1) = 0 . \tag{5}$$

This determines three sets of eigenvalues, $\{x_j\}$, $\{y_j\}$ and $\{z_j\}$, and three eigenfunctions, $c_j(t) = e^{-iz_j A(t)}$, where $A(t) = \int_0^t V(t')dt'$. Specifically, $c_j = \sum_{i=1}^3 \mathcal{M}_{ji}a_i$, where,

$$\mathcal{M} = \begin{pmatrix} 1 & x_1 & y_1 \\ 1 & x_2 & y_2 \\ 1 & x_3 & y_3 \end{pmatrix} \tag{6}$$

This matrix, $\mathcal{M}$, may be inverted, namely,

$$\mathcal{M}^{-1} = \frac{1}{\Delta} \begin{pmatrix} x_2y_3 - x_3y_2 & x_3y_1 - x_1y_3 & x_1y_2 - x_2y_1 \\ y_2 - y_3 & y_3 - y_1 & y_1 - y_2 \\ x_3 - x_2 & x_1 - x_3 & x_2 - x_1 \end{pmatrix} \tag{7}$$

where $\Delta = \det(\mathcal{M}) = x_1y_2 + x_2y_3 + x_3y_1 - x_1y_3 - x_2y_1 - x_3y_2$.

Now one may express the unperturbed state amplitudes, $a_j(t)$, in terms of the dressed-state amplitudes, $c_j(t)$. Using $c_j(t) = e^{-iz_j A(t)}$, one has $a_i(t) = \sum_{j=1}^3 \mathcal{M}_{ij}^{-1} c_j(t) =$

$\sum_{j=1}^3 \mathcal{M}_{ij}^{-1} e^{-iz_j A(t)}$. This leads to,

$$\begin{aligned}
P_1(t) &= |a_1(t)|^2 = \frac{1}{\Delta^2} [(x_2 y_3 - x_3 y_2)^2 + (x_3 y_1 - x_1 y_3)^2 + (x_1 y_2 - x_2 y_1)^2 \\
&\quad + 2(x_2 y_3 - x_3 y_2)(x_3 y_1 - x_1 y_3) \cos((z_1 - z_2)A(t)) \\
&\quad + 2(x_2 y_3 - x_3 y_2)(x_1 y_2 - x_2 y_1) \cos((z_1 - z_3)A(t)) \\
&\quad + 2(x_3 y_1 - x_1 y_3)(x_1 y_2 - x_2 y_1) \cos((z_2 - z_3)A(t))] , \\
P_2(t) &= |a_2(t)|^2 = \frac{1}{\Delta^2} [(y_2 - y_3)^2 + (y_3 - y_1)^2 + (y_1 - y_2)^2 \\
&\quad + 2(y_2 - y_3)(y_3 - y_1) \cos((z_1 - z_2)A(t)) \\
&\quad + 2(y_2 - y_3)(y_1 - y_2) \cos((z_1 - z_3)A(t)) \\
&\quad + 2(y_3 - y_1)(y_1 - y_2) \cos((z_2 - z_3)A(t))] , \\
P_3(t) &= |a_3(t)|^2 = \frac{1}{\Delta^2} [(x_3 - x_2)^2 + (x_1 - x_3)^2 + (x_2 - x_1)^2 \\
&\quad + 2(x_3 - x_2)(x_1 - x_3) \cos((z_1 - z_2)A(t)) \\
&\quad + 2(x_3 - x_2)(x_2 - x_1) \cos((z_1 - z_3)A(t)) \\
&\quad + 2(x_1 - x_3)(x_2 - x_1) \cos((z_2 - z_3)A(t))] . \tag{8}
\end{aligned}$$

Eq(8) generally gives the transition probabilities to all three final states for arbitrary $V_{jk}(t)$, and corresponds to the more general expression [26], $P_k(t) = \sum_i^n \sum_j^n \mathcal{M}_{ki}^{-1} \mathcal{M}_{kj}^{-1} \cos[(z_i - z_j)A(t)]$. Since the x_j 's, y_j 's and z_j 's vary with the $V_{jk}(t)$, one may then seek conditions on the matrix elements V_{jk} such that the electron populations $P_k(t_0)$ take any desired values at any specified time, $t = t_0$.

B. Complete population transfer

Next we seek conditions on the $V_{jk}(t)$ such that at time t_0 the electron population is completely transferred from its initial state $i = 1$ to a different final state, $i = 2$ or 3 . As shown from Eq(A7) in an appendix, the condition that $\epsilon_j = 0$ yields $\alpha = 2/y - y$ and

$\beta = \pm 1$. From the appendix taking $\beta = +1$ in case i) and using $r = \pm \sqrt{\frac{2}{n_1 n_2}}$, one has,

$$\begin{aligned}
\{x_1, x_2, x_3\} &= \{1, 1, -1\} \quad , \\
\{y_1, y_2, y_3\} &= \{y_+, y_-, 0\} = \frac{1}{2}\{-\alpha + \sqrt{\alpha^2 + 8}, -\alpha - \sqrt{\alpha^2 + 8}, 0\} \\
&= r \{n_1, -n_2, 0\} \quad , \\
\{z_1, z_2, z_3\} &= \{\alpha + y_+, \alpha + y_-, -\alpha\} = \frac{1}{2}\{\alpha + \sqrt{\alpha^2 + 8}, \alpha - \sqrt{\alpha^2 + 8}, -2\alpha\} \\
&= \{-y_-, -y_+, -\alpha\} = r \{n_2, -n_1, n_1 - n_2\} \quad , \\
\Delta &= 2(y_- - y_+) = -2\sqrt{\alpha^2 + 8} = -2r(n_1 + n_2) \quad .
\end{aligned} \tag{9}$$

Here n_1 and n_2 are integers whose values are specified below. This leads to certain allowed values of the action integral $A(t_0)$ and the relative interaction strength, α , namely,

$$\begin{aligned}
3r A(t_0) &= \pm 3\sqrt{\frac{2}{n_1 n_2}} A(t_0) = \pi \quad , \\
\alpha &= V_{12}(t)/V_{23}(t) = r (n_2 - n_1) \quad , \\
\beta &= V_{13}(t)/V_{23}(t) = \pm 1 \quad .
\end{aligned} \tag{10}$$

The same allowed values of $A(t_0)$, α and β are found in all cases, shown in the appendix.

One may use the values of the x_j , y_j , z_j and Δ from Eq(9) in Eq(8) to obtain explicit expressions for the transition probabilities, namely,

$$\begin{aligned}
P_1(t) &= \frac{1}{2(n_1 + n_2)^2} \{n_1^2 + n_2^2 + n_1 n_2 [1 + \cos((n_1 + n_2)rA(t))] \\
&\quad + (n_1 + n_2)[n_1 \cos((2n_1 - n_2)rA(t)) + n_2 \cos((2n_2 - n_1)rA(t))]\} \quad , \\
P_2(t) &= \frac{1}{2(n_1 + n_2)^2} \{n_1^2 + n_2^2 + n_1 n_2 [1 + \cos((n_1 + n_2)rA(t))] \\
&\quad - (n_1 + n_2)[n_1 \cos((2n_1 - n_2)rA(t)) + n_2 \cos((2n_2 - n_1)rA(t))]\} \quad , \\
P_3(t) &= \frac{2n_1 n_2}{(n_1 + n_2)^2} \sin^2 \left[\left(\frac{n_1 + n_2}{2} \right) rA(t) \right] \quad ,
\end{aligned} \tag{11}$$

where n_1, n_2 are odd integers. The constraints on the values of n_1 and n_2 are as follows. The condition $P_1 = 0$ requires that $\frac{1}{3}(2n_1 - n_2)$ and $\frac{1}{3}(2n_2 - n_1)$ are both odd integers (i.e. n_o and n'_o), and also requires that $\frac{1}{3}(n_1 + n_2)$ is an even integer. The condition $P_2 = 1$ imposes the same requirements. The condition $P_3 = 0$ only requires that $\frac{1}{3}(n_1 + n_2)$ be an even integer, but $P_3 \geq 0$ requires that $n_1 n_2 \geq 0$.

Finally a slightly neater solution may now be obtained from the requirements above that $2n_1 - n_2 = 3n_o$ and $2n_2 - n_1 = 3n'_o$, where n_o and n'_o are both odd integers (either positive

or negative). It is easily shown that $n_1 + n_2 = 3(n_o + n'_o)$ and $n_1 n_2 = 2n_o^2 + 5n_o n'_o + 2n_o'^2$. Then $r = \pm(n_o^2 + \frac{5}{2}n_o n'_o + n_o'^2)^{-\frac{1}{2}}$, $\alpha = r(n'_o - n_o)$ and $3rA(t_0) = \pi$. Now,

$$\begin{aligned} P_1(t) &= \frac{1}{2(n_1 + n_2)^2} \{n_1^2 + n_2^2 + n_1 n_2 [1 + \cos((n_o + n'_o)\pi (A(t)/A(t_0)))] \\ &\quad + (n_1 + n_2)[n_1 \cos(n_o \pi (A(t)/A(t_0))) + n_2 \cos(n'_o \pi (A(t)/A(t_0)))]\} , \\ P_2(t) &= \frac{1}{2(n_1 + n_2)^2} \{n_1^2 + n_2^2 + n_1 n_2 [1 + \cos((n_o + n'_o)\pi (A(t)/A(t_0)))] \\ &\quad - (n_1 + n_2)[n_1 \cos(n_o \pi (A(t)/A(t_0))) + n_2 \cos(n'_o \pi (A(t)/A(t_0)))]\} , \\ P_3(t) &= \frac{2n_1 n_2}{(n_1 + n_2)^2} \sin^2\left(\frac{1}{2}(n_o + n'_o)\pi (A(t)/A(t_0))\right) . \end{aligned} \quad (12)$$

Here n_o and n'_o are arbitrary odd integers, $n_1 = 2n_o + n'_o$, and $n_2 = n_o + 2n'_o$. At $t = t_0$, $A(t)/A(t_0) = 1$, so that $P_2(t_0) = 1$ with $P_1(t_0) = P_3(t_0) = 0$. This yields complete population transfer from level 1 to level 2 at $t = t_0$. We regard n_o and n'_o as the more fundamental numbers since they obey the simplest rules.

III. RESULTS

Some allowed values of the action integral, $A(t_0)$, and the relative interaction strength, α , are given in table I. This table includes values of $\{n_1, n_2\}$, $\{n_o, n'_o\}$ and all three $\{k, k'\}$ cases defined in the appendix. From more complete numerical output we confirm that n_1 and n_2 each acquire all possible odd integer values, although values of the product $n_1 \cdot n_2$ are restricted. This is consistent with the condition that $n_1 = 2n_o + n'_o$, and $n_2 = n_o + 2n'_o$, where n_o and n'_o are arbitrary odd integers. It is also evident that the even integer, $n_e = n_o + n'_o$, takes on all even values. One may also show algebraically that for each value of the even integer k in case i) there are two odd values of an odd integer k in case ii), and vice versa. The sets of integers $\{n_1, n_2\}$, $\{n_o, n'_o\}$ and $\{k, k'\}$ are redundant. The three sets of $\{k, k'\}$ correspond to a single set $\{n_1, n_2\}$, while the $\{n_o, n'_o\}$ are in one to one correspondence with the $\{n_1, n_2\}$.

Numerical calculations for the time dependence of the populations in a degenerate 3-state atom perturbed by external interactions with $V(t) = V_0 \cos(\omega t)$ are presented in figures 1-4. These results were obtained by using a standard fourth order Runge-Kutta numerical integration of Eq(3) with $\epsilon_j = 0$ and $\beta = 1$ for various values of α and $A(t_0)$. For most values of α the population transfer into either level 2 or level 3 is always incomplete. A typical case is shown in figure 1 where $\alpha = 2$ and $A(t_0) = 1.5$. We note that $P_1(t)$ does return to

$n_1 \cdot n_2$	n_1	n_2	n_e	n_o	n_o	n_o'	n_o'	n_e	$A(t_0)$	α
			k	$-k'$	k	k'	$-k$	k'		
5	± 1	± 5	± 2	∓ 1	∓ 1	± 3	± 3	± 2	± 1.656	∓ 2.530
5	± 5	± 1	± 2	± 3	± 3	∓ 1	∓ 1	± 2	± 1.656	± 2.530
9	± 3	± 3	± 2	± 1	± 1	± 1	± 1	± 2	± 2.221	0.000
11	± 1	± 11	± 4	∓ 3	∓ 3	± 7	± 7	± 4	± 2.456	∓ 4.264
11	± 11	± 1	± 4	± 7	± 7	∓ 3	∓ 3	± 4	± 2.456	± 4.264
17	± 1	± 17	± 6	∓ 5	∓ 5	± 11	± 11	± 6	± 3.053	∓ 5.488
17	± 17	± 1	± 6	± 11	± 11	∓ 5	∓ 5	± 6	± 3.053	± 5.488
23	± 1	± 23	± 8	∓ 7	∓ 7	± 15	± 15	± 8	± 3.551	∓ 6.487
23	± 23	± 1	± 8	± 15	± 15	∓ 7	∓ 7	± 8	± 3.551	± 6.487
27	± 3	± 9	± 4	∓ 1	∓ 1	± 5	± 5	± 4	± 3.848	∓ 1.633
27	± 9	± 3	± 4	± 5	± 5	∓ 1	∓ 1	± 4	± 3.848	± 1.633
29	± 1	± 29	± 10	∓ 9	∓ 9	± 19	± 19	± 10	± 3.988	∓ 7.353
29	± 29	± 1	± 10	± 19	± 19	∓ 9	∓ 9	± 10	± 3.988	± 7.353
35	± 1	± 35	± 12	∓ 11	∓ 11	± 23	± 23	± 12	± 4.381	∓ 8.128
35	± 5	± 7	± 4	± 1	± 1	± 3	± 3	± 4	± 4.381	∓ 0.478
35	± 7	± 5	± 4	± 3	± 3	± 1	± 1	± 4	± 4.381	± 0.478
35	± 35	± 1	± 12	± 23	± 23	∓ 11	∓ 11	± 12	± 4.381	± 8.128

TABLE I: Some values for the action integral, $A(t_0)$, and the relative interaction strength, α , allowed for total population transfer. These values are found using Eq(10) subject to the conditions listed. Here $n_e = n_o + n'_o$, $n_o = \frac{1}{3}(2n_1 - n_2)$ and $n'_o = \frac{1}{3}(2n_2 - n_1)$.

zero twice in each period, T , of the external field, but that complete population transfer to an initially unoccupied state never occurs. We also ran calculations (not shown) for $\alpha = 8.5$ and $A(t_0) = 5.8$, which are not values that lead to complete population transfer from Eq(10). Again complete transfer to an initially unoccupied level never occurred. However, there were rapid oscillations in the populations of all states except near $t = T/4$ and $t = 3T/4$, where none of the populations oscillated rapidly. This was similar to figure 4. This appears to correspond to the onset of complete population transfer, which occurs at $t = T/4$ and odd multiples of $T/4$, as seen in figures 2-4.




FIG. 1: Occupation probabilities as a function of time over one period, T , of the external field, $V(t) = V_0 \cos(\omega t)$. The solid line is the probability, $P_2(t)$, that the electron is transferred to the target state, namely state 2. The long dash line is the probability, $P_1(t)$, that the electron is in state 1, the state in which it began initially. The short dash line is the probability, $P_3(t)$, that the electron is in state 3. In this figure $\alpha = V_{12}/V_{23} = 2$ and the action area at $t = t_0$ is $A(t_0) = \int_0^{t_0} V(t') dt' = 1.5$. Since α and $A(t_0)$ do not correspond to values given by Eq(10), complete transfer to level 2 does not occur.




FIG. 2: Occupation probabilities as a function of time. The solid line denotes $P_2(t)$; the long dash line denotes $P_1(t)$; and the short dash line denotes $P_3(t)$. In this figure we use the allowed values, $\alpha = V_{12}/V_{23} = 0$ and $A(t_0) = \int_0^{t_0} V(t') dt' = 2.221$, corresponding to $n_1 = 3$ and $n_2 = 3$ ($n_o = n'_o = 1$) in Eq(10). Complete transfer to level 2 from level 1 occurs at $t = t_0 = T/4$ and again at odd multiples of t_0 .




FIG. 3: Occupation probabilities as a function of time. The solid line denotes $P_2(t)$; the long dash line denotes $P_1(t)$; and short dash line denotes $P_3(t)$. In this figure $\alpha = -2.530$ and $A(t_0) = 1.656$, corresponding to $n_1 = 1$ and $n_2 = 5$ ($n_o = -1$, $n'_o = 3$) in Eq(10).

Calculations using a few values of α that permit complete transfer to level 2 are shown in figures 2-4. Our numerical codes give the same results as the analytic expressions of Eq(11). This provides a check that our algebra is correct. We note that $\frac{2n_1n_2}{(n_1+n_2)^2} \leq \frac{1}{2}$ and that the maximum value of $P_3(t)$ occurs for $n_1 = n_2$, where $P_{3 \max} = \frac{1}{2}$, consistent with figure 2. This corresponds to $\alpha = 0$ so that direct transitions from level 1 to level 2 are forbidden. Transfer to level 2 occurs via the intermediate level 3. Transfer from level 1 to level 2 and back is complete, and occurs periodically. In general $\alpha = 0$ occurs when $n_1 = n_2 = 3n_{\text{odd}}$, where n_{odd} is any odd integer. This appears to give the simplest condition that allows complete population transfer. In this case the action area is $A(t_0) = n_{\text{odd}}\pi/\sqrt{2}$.

Calculations for two other values of α that allow complete transfer to level 2 are shown in figures 3 and 4. We see that complete transfer occurs twice in one period of the oscillating field but that the frequency of the "side bands" increases as αA increases. We note that $\alpha = \pm\sqrt{\frac{2}{n_1n_2}}(n_2 - n_1)$ becomes either large ($n_2 \gg n_1$, or vice versa), or small ($n_1 \sim n_2$) as n_1n_2 increases, while $A(t_0)$ increases as $\sqrt{n_2n_2/2}$. When complete population transfer occurs, the population lingers in level 2, as seen in figures 2-4. It can be shown [25] that near $t = t_0$ $1 - P_2(t)$ varies as $[\omega(t - t_0)]^4$.

Additional control [25] may be achieved by changing the shape of $V(t)$. This can be used to control how long the population remains near unity in level 2, for example. An interesting example is the case of an ideal, sudden 'kick' produced by $V(t) = A_0\delta(t - t_0)$. If




FIG. 4: Occupation probabilities as a function of time. The solid line shows $P_2(t)$; the long dash line shows $P_1(t)$; and the short dash line shows $P_3(t)$. Here $\alpha = 8.128$ and $A(t_0) = 4.381$, corresponding to $n_1 = 35$ and $n_2 = 1$ ($n_o = 23$, $n'_o = -11$) in Eq(10). Since these values are allowed, the electron population is completely transferred to level 2 from level 1.

$A_0 \rightarrow n_{odd}\pi/\sqrt{2}$ for example, then the mere presence of level 3 allows transfer from level 1 to level 2 without any direct transfer from level 1 to level 2. With this ideal 'kick', level 2 is unoccupied before $t = t_0$, and fully occupied after $t = t_0$.

To obtain analytic solutions to Eq.(2) for a 3-level atom, we imposed degeneracy on the 3-level manifold, e.g. by taking $E_j = E = 0$. If $E \neq 0$, then E can be removed from Eq(1) by an overall phase transformation, e^{iEt} . When the levels are not degenerate $\omega_{ij} = (E_i - E_j)/\hbar \neq 0$, and the population transfer is incomplete. The effect of this population leakage can be calculated by expanding the transition amplitude a_2 in a power series in time with terms small in $\omega_{ij}t$. After some algebra we obtain from Eq(1) the difference between the degenerate and nearly degenerate populations of level 2, namely, $\Delta P_2(t) \simeq \frac{1}{12}[2(2\omega_{13} - \omega_{12})V_{12}(0)V_{13}(0)V_{23}(0) + \omega_{12}^2 V_{12}^2(0)]t^4$, where $V_{ij}(0)$ denotes the value of $V_{ij}(t)$ at $t = 0$. By the time the occupation probability P_2 reaches its first maximum at $t_0 = T/4 = \frac{\pi}{2\omega}$, using $\alpha^{-1}V_{12}(0) = \beta^{-1}V_{13}(0) = V_{23}(0) = V(0) = \pm\sqrt{\frac{n_1 n_2}{2}}\frac{\pi}{3}\omega$, the difference becomes, $\Delta P_2(t_0) \simeq \frac{1}{27}(\frac{\pi}{2})^6[\frac{\pi}{3}\beta n_1 n_2 (n_2 - n_1)(2\frac{\omega_{13}}{\omega} - \frac{\omega_{12}}{\omega}) + (n_2 - n_1)^2(\frac{\omega_{12}}{\omega})^2]$. The choice of the field frequency, ω , involves a trade-off between the duration of time the population remains in state 2 and the population leakage. The higher the frequency, ω , the smaller the population leakage, ΔP_2 , but the shorter the duration time, T_s , that the population

remains in level 2. This effect of population leakage is similar as that for a 2-level atom [25]. Numerical calculations indicate that this population leakage grows rapidly in time.

IV. DISCUSSION

In this paper we have considered transitions amplitudes and probabilities in a 3-level atom with degenerate states. We also assumed that the interaction matrix elements, $V_{ij}(t)$ all have a common time dependence. In realistic atomic systems, energy levels are seldom, if ever, exactly degenerate. Also levels outside 3-level manifold usually exist. Consequently, use of our results is restricted to external fields with frequencies in the range, $\omega_{min} < \omega < \omega_{max}$. Here $\hbar\omega_{min}$ is the energy splitting of the nearly degenerate states, and $\hbar\omega_{max}$ is the energy difference between the 3-level manifold and the closest state in energy outside the manifold.

It is instructive to compare our 3-level results to simpler 2-level results [25, 26]. The equations for a 2-level atom may be recovered from our 3-level equations by taking V_{12} and V_{13} to zero, which corresponds to $\alpha \rightarrow \infty$ (with any $|\epsilon_j/\alpha| < \infty$), as may be understood from Eq(4). For the 2-level atom it has been shown [25, 26] that $a_j(t)$ may be expressed in terms of eigenstates $c_i(t)$ via the relation, $c_i(t) = \sum_{j=1}^2 \mathcal{M}_{ij} a_j(t)$, where $\mathcal{M}_{ij} = \begin{pmatrix} 1 & y_+ \\ 1 & y_- \end{pmatrix}$. Here $y_{\pm}$ are the same as those given by Eq(A8). This matrix may be inverted to give the population amplitudes, $a_1(t) = \frac{1}{\Delta}[y_- c_1(t) - y_+ c_2(t)]$ and $a_2(t) = \frac{1}{\Delta}[-c_1(t) + c_2(t)]$, where $\Delta = 2\sqrt{1 + (\frac{\epsilon_2 - \epsilon_1}{2})^2}$ is the determinant of $\mathcal{M}_{ij}$. A simple solution occurs when $\epsilon_1 = \epsilon_2 = \epsilon$, namely, $a_1(t) = e^{-i\epsilon A(t)} \frac{1}{2}[e^{-iA(t)} + e^{+iA(t)}] = e^{-i\epsilon A(t)} \cos(A(t))$ and $a_2(t) = e^{-i\epsilon A(t)} \frac{1}{2}[e^{-iA(t)} - e^{+iA(t)}] = -ie^{-i\epsilon A(t)} \sin(A(t))$. Note that $P_1 = 1 - P_2$. One may then simply determine the conditions under which the population of level 2, $P_2(t_0)$, takes on any desired value at time t_0 . If $\epsilon_1 \neq \epsilon_2$, it is easily shown that $P_2(t_0) \leq 1/(1 + (\epsilon_2 - \epsilon_1)^2/4) < 1$. Thus the diagonal matrix elements of $V_{ij}(t)$ prevent complete transfer to the initially unoccupied level when ϵ_1 and ϵ_2 are unequal, and, when they are equal, they simply contribute an overall phase. When $\epsilon_1 = \epsilon_2$ any value of $P_2(t_0)$ between 0 and 1 can be found. In particular, $P_2(t_0) = 1$ if $A(t_0) = \int_0^{t_0} V_{12}(t') dt' = n_{odd}\pi/2$. This allowed value of $A(t_0)$ for complete transfer to level 2 differs from the allowed values given in Eq(10) for degenerate 3-level atoms. By comparison to a 2-level atom the conditions for complete transfer are generally more complex in a 3-level atom, as discussed in the appendix. For example, complete

population transfer can occur in a 3-level atom when $\epsilon_2 \neq \epsilon_3$. In both cases, however, the duration of time spent in the transferred state can be controlled by adjusting the time dependence of the external field. Also if the states are not quite degenerate then population leakage occurs. For a harmonic $V(t)$ the population leakage in a 2-level atom varies as $\Delta P_2(t_0) \simeq \frac{1}{4}(\frac{\pi}{2})^6(\omega_{12}/\omega)^2$. For $2s - 2p$ transitions in atomic hydrogen complete population control can be nearly achieved in this manner using a radiation field with wavelengths (and intensities) ranging from a few μm (with about 10^{12} W/cm²) up to a few cm (with about 10^4 W/cm²). At wavelengths below a few μm coupling to nearby $n = 3$ atomic levels can be significant, and this 2-level model breaks down.

While we have not provided calculations in this paper for specific experiments, some general guidelines for experimental tests and applications are evident. First, the model must be valid, so that the frequency of the external interaction is limited by $\omega_{min} < \omega < \omega_{max}$. If the external interaction varies harmonically, $V(t) = V_0 \cos(\omega t)$, then the first line of Eq(10) imposes a constraint between V_0 and ω , namely, $V_0/\omega = \pm \sqrt{\frac{n_1 n_2}{2}} \frac{\pi}{3}$. For a harmonic interaction the duration of the time, T_s , that the population remains in level 2 varies inversely with ω ; the higher the frequency, ω , the smaller the time the population remains in level 2. Our degenerate 3-level model can be applied to systems with dipole selection rules. A dipole selection rule can correspond to $\alpha = 0$. Transfer from level 1 to level 2 (which is the dipole forbidden transition) is complete at time t_0 if the two allowed transition matrix elements, V_{23} and V_{13} , are equal in absolute magnitude and $A(t_0) = n_{odd}\pi/2$. In some atomic systems state 2 could decay, with a lifetime, T_d , to another level outside the degenerate manifold and be lost. Such loss can be controlled by adjusting ω . Our model also rests on degeneracy. If the three levels are not degenerate the transfer of population to level 2 is incomplete. We call this population leakage. As discussed above this population leakage may be minimized by using high frequency external interactions. i.e. $\omega \gg \omega_{min}$, but at a cost to duration of time, T_s , the population remains in level 2.

Using nearly degenerate states, the duration time, T_s , can be further controlled [25] by adjusting the shape of $V(t)$. As mentioned in section III for example, the transfer is complete, instantaneous and permanent when $V(t) = n_{odd} \frac{\pi}{\sqrt{2}} \delta(t - t_0)$. Such a quick, hard pulse is called [27, 28, 29] a 'kick'. Two practical limitations on this ideal model are the impossibility of producing a signal that varies as $\delta(t - t_0)$, and the existence of an infinitely wide spectrum of high frequency components with frequencies $\omega > \omega_{max}$ in the Fourier

spectrum of $\delta(t - t_0)$. Fortunately these two difficulties can both be addressed by using 'kicks' of finite width in time. In some cases it may be possible to design a 'kick' so that its duration is short compared to any other changes in the system, so that a finite 'kick' may be sensibly represented by $\delta(t - t_0)$. If, in addition, the energy levels outside the (nearly) degenerate manifold have a large energy gap $\hbar\omega_{max}$, then it may be possible that $\omega_{min} < \omega < \omega_{max}$ and our model may be applicable. Applying $V(t) \simeq \delta(t - t_0)$ to nearly degenerate atomic systems leads to the 'gedanken' question, what happens when one tries to force the transition to occur within a small time interval about t_0 in a degenerate quantum system where Δt is large?

Let us next consider some mathematical features that occur when degenerate states are used. The exact probability amplitude for a transition from an initial state $|i\rangle$ to a final state $|f\rangle$ may be expressed in terms of the time evolution operator $U(t_i, t_f)$, namely, $a_{fi} = \langle f|U(t_f, t_i)|i\rangle$, where (setting $t_i = 0$ and $t_f = t$), $U(t, 0) = T e^{-i \int_0^t V(t') dt'} = \sum_n \frac{(-i)^n}{n!} \int_0^t \dots \int_0^t T V(t_n) \dots V(t_1) dt_n \dots dt_1$. Here T is the Dyson time ordering operator which places the interactions in the order of increasing time so that for $t \geq t'$, $TV(t)V(t')$ and $TV(t)V(t') = 0$, and vice versa for $t' \geq t$, whence $T[V(t), V(t')] \neq 0$. In any intermediate time step from t_j to t_{j+1} , the matrix element is $\int_0^t e^{(E_I - E_{I'})t_j} < I|V(t_j)|I' > dt_j$. Without the phase $e^{(E_I - E_{I'})t_j}$ there is no time ordering since $T[V(t), V(t')] \rightarrow 0$ as $T \rightarrow 1$. If all the states are degenerate, then $\Delta E = E_I - E_{I'} \rightarrow 0$ in every time step. This corresponds to taking $T \rightarrow 1$ in $U(t, 0) = T e^{-i \int_0^t V(t') dt'}$. In this limit of degeneracy, $U(t, 0) \rightarrow e^{-i \int_0^t V(t') dt'}$, corresponding to the approximation of Magnus [30], where $[V(t), V(t')] \rightarrow 0$. In this limit there are no time correlations in the time propagation. Imposing degeneracy removes time ordering. Moreover in standard stationary scattering with $H = H_0 + V$, the Fourier transform of $e^{i \int_0^t (E - H) dt}$ is $i\pi\delta(E - H)$, which is the energy conserving part of the Green function, $G_0 = \frac{1}{E - H \pm i\eta} = i\pi\delta(E - H) \mp P_v \frac{1}{E - H}$. The energy non-conserving quantum fluctuations, $P_v \frac{1}{E - H}$, imposed by the asymptotic $\pm i\eta$ initial condition, carry the direction of time, i.e., outgoing or incoming scattered waves. Here $P_v \frac{1}{E - H}$ is the principal value integral that omits the contribution at $E = H$. The Fourier transform of $P_v \frac{1}{E - H}$ is $\text{sign}(t) = T - 1$. Degeneracy, $P_v \frac{1}{E - H} \rightarrow 0$, again removes time ordering. Thus, imposing degeneracy mathematically

corresponds to,

$$\begin{aligned}
\Delta E \rightarrow 0 &\iff \int_0^t e^{(E_I - E_{I'})t_j} < I|V(t_j)|I' > dt_j \rightarrow \int_0^t < I|V(t_j)|I' > dt_j \\
&\iff T[V(t), V(t')] \rightarrow 0 \iff U(t, 0) \rightarrow e^{-i \int_0^t V(t') dt'} \\
&\iff e^{i(E-H_0)t} T e^{-i \int_0^t V(t') dt'} \rightarrow e^{i \int_0^t (E-H) dt} \\
&\iff \frac{1}{E - H \pm i\eta} \rightarrow \frac{1}{E - H} \iff P_v \frac{1}{E - H} \rightarrow 0 \\
&\iff T \rightarrow 1 \quad .
\end{aligned} \tag{13}$$

As a consequence of using degenerate states, quantum energy fluctuations in intermediate states are eliminated. The Hilbert space is restricted so that $\Delta t = \hbar/\Delta E \rightarrow \infty$. In addition the minimal size of the wavepacket, $\Delta \ell = v\Delta t$, now extends to infinity, so that the quantum system may not be decoherently decoupled from its macroscopic environment. This effect is evident in $2s - 2p$ transitions in hydrogen caused by the impact of ions, where the range of the interaction goes to infinity as the levels become degenerate [31, 32]. Within the interval Δt time cannot be localized, e.g. reproducibly observed. Time ordering i.e. causality, is not operative and there is no 'flow of time'. In the classical limit as $\Delta t \rightarrow 0$, time can be measurably localized, being defined by two closely spaced decoherent observable events. In this classical limit causality strictly holds.

We note that at sufficiently high ω all bound levels in the atomic elements become nearly degenerate. Hence, if one can deal with high energy continuum states, our approach might be useful in applications involving fourth generation synchrotrons that produce intense high frequency fields. An extension of this method might also be used for control transitions in high Rydberg states [33], including adiabatic rapid passage [34, 35]. RWA is used to describe coherent storage of information in photonic states [13]. The approach developed here may be useful for modeling information transmission and storage in atomic states [10] in a new way.

V. SUMMARY

We have shown that in a 3-level atom with degenerate energies, electron population is completely transferred via an external interaction at a designated time, t_0 , from an initially occupied level (level 1) to a designated initially unoccupied level (level 2) under two

conditions. The first condition for complete transfer is that the ratio of the matrix elements of the external interaction, $V_{ij}(t)$, satisfy $V_{12}(t)/V_{23}(t) = \alpha = \pm \sqrt{\frac{2}{n_1 n_2}}(n_2 - n_1)$, and $V_{13}(t)/V_{23}(t) = \beta = \pm 1$. The second condition is that at $t = t_0$ the action area of $V(t)$ satisfy $A(t_0) = \int_0^{t_0} V(t') dt' = \pm \sqrt{\frac{n_1 n_2}{2}} \frac{\pi}{3}$, where we have set $V_{23}(t) = V(t)$. Here n_1 and n_2 are integers such that $n_1 = 2n_o + n'_o$ and $n_2 = n_o + 2n'_o$, where n_o and n'_o are any odd integers. The duration of time the transferred population remains in level 2 can be controlled either by varying the frequency, ω , of the external potential, or by varying the shape of $V(t)$.

Acknowledgments

We thank J.H. Eberly, B.W. Shore and A.L. Godunov for useful discussion. This work was supported in part by the Division of Chemical Sciences, Office of Sciences, U.S. Department of Energy. KhR is supported by a NSF-NATO Fellowship.

APPENDIX A: METHOD TO FIND CONDITIONS FOR COMPLETE POPULATION TRANSFER

Here we show how to find the conditions on the V_{jk} such that the electron population is completely transferred from its initial state $i = 1$ to a different final state, $i = 2$ or 3 . A convenient way to begin is to take $dP_k(t)/dt = 0$ at $t = t_0$. This gives both maxima and minima for P_k . Using Eq(8) this yields,

$$\sin((z_i - z_j)A(t_0)) = 0 \quad (\text{A1})$$

$$(z_i - z_j)A(t_0) = m_{ij}\pi \quad (m_{ij} \neq 0) \quad (i, j = 1, 2, 3) \quad .$$

Using $\cos((z_i - z_j)A(t_0)) = \cos(m_{ij}\pi) = (-1)^{m_{ij}}$, one has,

$$\begin{aligned} P_1 &= |a_1|^2 = \frac{1}{\Delta^2} [(x_2 y_3 - x_3 y_2)^2 + (x_3 y_1 - x_1 y_3)^2 + (x_1 y_2 - x_2 y_1)^2 \\ &\quad + 2(-1)^{m_{12}}(x_2 y_3 - x_3 y_2)(x_3 y_1 - x_1 y_3) \\ &\quad + 2(-1)^{m_{13}}(x_2 y_3 - x_3 y_2)(x_1 y_2 - x_2 y_1) \\ &\quad + 2(-1)^{m_{23}}(x_3 y_1 - x_1 y_3)(x_1 y_2 - x_2 y_1)] \\ &= \frac{1}{\Delta^2} [(-1)^{k_1}(x_2 y_3 - x_3 y_2) + (-1)^{k_2}(x_3 y_1 - x_1 y_3) + (-1)^{k_3}(x_1 y_2 - x_2 y_1)]^2 \quad , \quad (\text{A2}) \end{aligned}$$

where $k_1 + k_2 = m_{12}$, $k_1 + k_3 = m_{13}$ and $k_2 + k_3 = m_{23}$. Now,

$$\begin{aligned}
k &\equiv k_1 = \frac{1}{2}(m_{12} + m_{13} - m_{23}) = \frac{1}{2}\left(\frac{A(t_0)}{\pi}(z_1 - z_2) + \frac{A(t_0)}{\pi}(z_1 - z_3) - \frac{A(t_0)}{\pi}(z_2 - z_3)\right) \\
&= \frac{A(t_0)}{\pi}(z_1 - z_2) \\
k_2 &= \frac{1}{2}(m_{12} + m_{23} - m_{13}) = \frac{1}{2}\left(\frac{A(t_0)}{\pi}(z_1 - z_2) + \frac{A(t_0)}{\pi}(z_2 - z_3) - \frac{A(t_0)}{\pi}(z_1 - z_3)\right) = 0 \\
k' &\equiv k_3 = \frac{1}{2}(m_{13} + m_{23} - m_{12}) = \frac{1}{2}\left(\frac{A(t_0)}{\pi}(z_1 - z_3) + \frac{A(t_0)}{\pi}(z_2 - z_3) - \frac{A(t_0)}{\pi}(z_1 - z_2)\right) \\
&= \frac{A(t_0)}{\pi}(z_2 - z_3)
\end{aligned} \tag{A3}$$

This yields extrema for P_k at $t = t_0$, namely,

$$\begin{aligned}
P_1 &= \frac{1}{\Delta^2}[(-1)^k(x_2y_3 - x_3y_2) + (x_3y_1 - x_1y_3) + (-1)^{k'}(x_1y_2 - x_2y_1)]^2 \\
P_2 &= \frac{1}{\Delta^2}[(-1)^k(y_2 - y_3) + (y_3 - y_1) + (-1)^{k'}(y_1 - y_2)]^2 \\
P_3 &= \frac{1}{\Delta^2}[(-1)^k(x_3 - x_2) + (x_1 - x_3) + (-1)^{k'}(x_2 - x_1)]^2
\end{aligned} \tag{A4}$$

where k and k' are non zero integers that are *not both even*.

a. Conditions for complete transfer

We now seek conditions such that at $t = t_0$, $P_1 = 0$, $P_2 = 1$, and $P_3 = 0$. There are three sets, $\{k, k'\}$, that satisfy these conditions, namely, i) k is even and k' is odd, ii) k and k' are both odd, and iii) k is odd and k' is even.

Case i: k is even and k' is odd. Then,

$$\begin{aligned}
P_1 &= \frac{1}{\Delta^2}[(x_2y_3 - x_3y_2) + (x_3y_1 - x_1y_3) - (x_1y_2 - x_2y_1)]^2 \\
P_2 &= \frac{1}{\Delta^2}[(y_2 - y_3) + (y_3 - y_1) - (y_1 - y_2)]^2 = \frac{4}{\Delta^2}(y_2 - y_1)^2 \\
P_3 &= \frac{1}{\Delta^2}[(x_3 - x_2) + (x_1 - x_3) - (x_2 - x_1)]^2 = \frac{4}{\Delta^2}(x_1 - x_2)^2 .
\end{aligned} \tag{A5}$$

From the condition that $P_2 = 1$ one has $[2(y_2 - y_1)/\Delta]^2 = 1$, where $y_1 \neq y_2$, so that $\Delta = \pm 2(y_2 - y_1)$. The condition that $P_3 = 0$ gives $[2(x_1 - x_2)/\Delta]^2 = 0$, which yields $x_1 = x_2 = x$. Using this result one then obtains $P_1 = [(x + x_3)(y_1 - y_2)/\Delta]^2$. Requiring $P_1 = 0$, one has $x_3 = -x$. Now, using $x_1 = x_2 = -x_3 = x$, we have from below Eq(7), $\Delta = 2x(y_2 - y_1)$. Finally since $\Delta = \pm 2(y_2 - y_1)$, one has $x_1 = x_2 = -x_3 = x = \pm 1$.

Case ii: k and k' are both odd. Then,

$$\begin{aligned} P_1 &= \frac{1}{\Delta^2} [-(x_2 y_3 - x_3 y_2) + (x_3 y_1 - x_1 y_3) - (x_1 y_2 - x_2 y_1)]^2 \\ P_2 &= \frac{1}{\Delta^2} [-(y_2 - y_3) + (y_3 - y_1) - (y_1 - y_2)]^2 = \frac{4}{\Delta^2} (y_3 - y_1)^2 \\ P_3 &= \frac{1}{\Delta^2} [-(x_3 - x_2) + (x_1 - x_3) - (x_2 - x_1)]^2 = \frac{4}{\Delta^2} (x_1 - x_3)^2. \end{aligned} \quad (\text{A6})$$

From the condition that $P_2 = 1$ one has $[2(y_3 - y_1)/\Delta]^2 = 1$, where $y_1 \neq y_3$, so that $\Delta = \pm 2(y_3 - y_1)$. The condition that $P_3 = 0$ gives $[2(x_1 - x_3)/\Delta]^2 = 0$, which yields $x_1 = x_3 = x$. Then one obtains $P_1 = [(x + x_2)(y_1 - y_3)/\Delta]^2$. When $P_1 = 0$ one has $x_2 = -x$. Now, using $x_1 = -x_2 = x_3 = x$, we get from below Eq(7), $\Delta = 2x(y_1 - y_3)$. Since $\Delta = \pm 2(y_3 - y_1)$, one has $x_1 = -x_2 = x_3 = x = \mp 1$.

Case iii: k is odd and k' is even. This is similar to case i). One obtains, $\Delta = 2x(y_3 - y_2)$ and $-x_1 = x_2 = x_3 = x = \pm 1$.

Complete population transfer, i.e. $P_2 = 1$, requires that $x = \pm 1$, so that the relative interaction strengths α and β must satisfy,

$$\begin{aligned} \alpha &= \frac{2}{y} - y \pm \frac{\epsilon_2 - \epsilon_1}{y^2} \pm (\epsilon_3 - \epsilon_2) \\ \beta &= \pm 1 + \frac{\epsilon_2 - \epsilon_1}{y} \end{aligned} \quad (\text{A7})$$

Next we use these results to find the conditions on V_{ij} required for complete population transfer to occur.

b. Quantum numbers for $A(t_0)$ and α

We now note that relatively simple conditions occur when $\epsilon_1 = \epsilon_2 = \epsilon_3 = \epsilon$, namely, $\alpha = \frac{2}{y} - y$ and $\beta = \pm 1$. In this case Eq(A7) reduces from a cubic to an easier quadratic equation in y . This quadratic equation corresponds to three roots of the more general cubic equation, namely,

$$\{y_j\} = \{y_+, y_-, 0\} = \left\{ \frac{-\alpha + \sqrt{\alpha^2 + 8}}{2}, \frac{-\alpha - \sqrt{\alpha^2 + 8}}{2}, 0 \right\}. \quad (\text{A8})$$

Now for case i) when k is even and k' is odd, one has from above, $x = \{\pm 1, \pm 1, \mp 1\}$ and $y = \{y_+, y_-, 0\}$. From Eq(A8), $z = \{\mp y_-, \mp y_+, \mp \alpha\}$. Then, with $E = A(t_0)/\pi$, the useful integers k and k' are: $k = \frac{A(t_0)}{\pi}(z_1 - z_2) = \pm E\sqrt{\alpha^2 + 8}$, and $k' = \frac{A(t_0)}{\pi}(z_2 - z_3) = \pm \frac{E}{2}(3\alpha - \sqrt{\alpha^2 + 8})$. After a little algebra one obtains $(k - k')(2k + k') = 18E^2 = n_1 n_2$.

For case ii) when k and k' are both odd, $x = \{\pm 1, \mp 1, \pm 1\}$ and $y = \{y_+, 0, y_-\}$. and $z = \{\mp y_-, \mp \alpha, \mp y_+\}$. Then, $k = \frac{A(t_0)}{\pi}(z_1 - z_2) = \pm \frac{E}{2}(3\alpha + \sqrt{\alpha^2 + 8})$, and $k' = \frac{A(t_0)}{\pi}(z_2 - z_3) = \mp \frac{E}{2}(3\alpha - \sqrt{\alpha^2 + 8})$. One may now verify that $(2k + k')(k + 2k') = 18E^2 = n_1 n_2$. For case iii) when k is odd and k' is even, $x = \{\mp 1, \pm 1, \pm 1\}$ and $y = \{0, y_+, y_-\}$, and $z = \{\mp \alpha, \mp y_-, \mp y_+\}$. Then, $k = \frac{A(t_0)}{\pi}(z_1 - z_2) = \mp \frac{E}{2}(3\alpha + \sqrt{\alpha^2 + 8})$, and $k' = \frac{A(t_0)}{\pi}(z_2 - z_3) = \pm E\sqrt{\alpha^2 + 8}$. One may again verify that $(2k' + k)(k' - k) = 18E^2 = n_1 n_2$.

Using $\alpha^2 = k^2/E^2 - 8$ one readily finds in case i), where k is even and k' is odd, that $k = \frac{1}{3}(n_1 + n_2)$ and $k' = \frac{1}{3}(n_2 - 2n_1)$. In case ii), where k and k' are both odd, one has $k = \frac{1}{3}(2n_1 - n_2)$ and $k' = \frac{1}{3}(2n_2 - n_1)$. In case iii) $k = \frac{1}{3}(n_1 - 2n_2)$ and $k' = \frac{1}{3}(n_1 + n_2)$. In all three cases one readily obtains the same equation for the allowed values of $A(t_0)$ and α , namely,

$$\begin{aligned} A(t_0) &= \pm \sqrt{\frac{n_1 n_2}{2}} \frac{\pi}{3} \\ \alpha &= \pm \sqrt{\frac{2}{n_1 n_2}} (n_2 - n_1) . \end{aligned} \quad (\text{A9})$$

It is remarkable that all three cases give the same formula for complete population transfer. The top equation corresponds to a constraint on the value of the action, $\int_0^t V(t') dt'$, i.e. the "area" of the external interaction, $V(t)$, or the phase shift caused by the external interaction. The bottom equation above gives the relative values of the V_{ij} .

APPENDIX B: GENERAL SOLUTIONS FOR THE DEGENERATE 3-LEVEL ATOM

In the main text a solution for the degenerate 3-level atom was found in the case when $\epsilon_j = 0$. We now choose $\epsilon_j^{-1} V_{jj}(t) = V(t) = V_{23}(t) = \alpha^{-1} V_{12}(t) = \beta^{-1} V_{13}(t)$. Then Eq(2) becomes,

$$\begin{aligned} i\dot{a}_1(t) &= \epsilon_1 V(t) a_1(t) + \alpha V(t) a_2(t) + \beta V(t) a_3(t) \\ i\dot{a}_2(t) &= \alpha V(t) a_1(t) + \epsilon_2 V(t) a_2(t) + V(t) a_3(t) \\ i\dot{a}_3(t) &= \beta V(t) a_1(t) + V(t) a_2(t) + \epsilon_3 V(t) a_3(t) . \end{aligned} \quad (\text{B1})$$

Consider a linear superposition $c(t) = a_1(t) + x a_2(t) + y a_3(t)$, where x and y are some

time-independent coefficients that we next determine. Then,

$$\begin{aligned}
i\dot{c}(t) &= i\dot{a}_1(t) + x i\dot{a}_2(t) + y i\dot{a}_3(t) \\
&= \epsilon_1 V(t) a_1(t) + \alpha V(t) a_2(t) + \beta V(t) a_3(t) \\
&\quad + x[\alpha V(t) a_1(t) + \epsilon_2 V(t) a_2(t) + V(t) a_3(t)] \\
&\quad + y[\beta V(t) a_1(t) + V(t) a_2(t) + \epsilon_3 V(t) a_3(t)] \quad .
\end{aligned} \tag{B2}$$

Set $z = \epsilon_1 + \alpha x + \beta y$. Then, $i\dot{c}(t) = zV(t)[a_1(t) + (\frac{\alpha + \epsilon_2 x + y}{\epsilon_1 + \alpha x + \beta y})a_2(t) + (\frac{\beta + x + \epsilon_3 y}{\epsilon_1 + \alpha x + \beta y})a_3(t)]$. We require that $i\dot{c}(t) = zV(t)c(t)$. This holds if and only if $z = \epsilon_1 + \alpha x + \beta y$, $x = (\alpha + \epsilon_2 x + y)/(\epsilon_1 + \alpha x + \beta y)$ and $y = (\beta + x + \epsilon_3 y)/(\epsilon_1 + \alpha x + \beta y)$. After some algebra this leads to the useful cubic equation,

$$\begin{aligned}
&[(\beta^2 - \alpha^2) + \alpha\beta(\epsilon_2 - \epsilon_3)]y^3 + [\alpha(2 - \alpha^2 - \beta^2) + \beta(2\epsilon_1 - \epsilon_2 - \epsilon_3) + \alpha(\epsilon_1 - \epsilon_3)(\epsilon_2 - \epsilon_3)]y^2 \\
&+ [(2\alpha^2 - \beta^2 - 1) + \alpha\beta(2\epsilon_3 - \epsilon_1 - \epsilon_2) + (\epsilon_1 - \epsilon_2)(\epsilon_1 - \epsilon_3)]y + [\alpha(\beta^2 - 1) - \beta(\epsilon_1 - \epsilon_2)] \\
&= 0 \quad .
\end{aligned} \tag{B3}$$

This determines three sets of eigenvalues, $\{x_j\}$, $\{y_j\}$ and $\{z_j\}$, and three eigenfunctions, $c_j(t) = e^{-iz_j A(t)}$, where $A(t) = \int_0^t V(t')dt'$.

When $\epsilon_j \neq 0$ complete population transfer can also occur. Using $\epsilon_j = \epsilon \neq 0$ adds nothing new, since it produces only an overall phase $e^{i\epsilon t}$. A slightly more general, but less elegant, solution for the degenerate 3-level equations is found when $\epsilon_1 = \epsilon_2 \neq \epsilon_3$. Again in this case $\alpha = \frac{2}{y} - y$ and $\beta = \pm 1$. The quadratic equation for y now becomes $y^2 + \frac{\alpha(1-\alpha^2) + \alpha(\epsilon_1 - \epsilon_3)^2 \pm (\epsilon_1 - \epsilon_3)}{1 - \alpha^2 \pm \alpha(\epsilon_1 - \epsilon_3)}y - 2 = 0$. Taking $\tilde{\alpha} = \frac{\alpha(1-\alpha^2) + \alpha(\epsilon_1 - \epsilon_3)^2 \pm (\epsilon_1 - \epsilon_3)}{1 - \alpha^2 \pm \alpha(\epsilon_1 - \epsilon_3)}$ the algebra is the same as that one above, except that $\alpha \rightarrow \tilde{\alpha}$. This includes an additional parameter, $(\epsilon_1 - \epsilon_3)$, gives simple allowed values of $\tilde{\alpha}$, and imposes a constraint between α and $(\epsilon_1 - \epsilon_3)$. More complex conditions for complete population transfer might exist when $\epsilon_1 \neq \epsilon_2$, in which case Eq(B3) is cubic in y . Finally it might be possible that solutions exist when the matrix elements $V_{ij}(t)$ have different time dependences, i.e. when α and β depend on t .

Conditions for complete transfer into level 3 at $t = t_0$ may be generally found similarly by interchange of the indices 2 and 3, with corresponding interchange of α with β and x with y .

[1] G. Alzetta, A. Gozzini, L. Moi, and G. Orriols, Nuovo Chimento Lett. **B 36**, 5 (1976).

- [2] A. Aspect, E. Arimondo, R. Kaiser, N. Vansteenkiste, and C. Cohen-Tannoudji, Phys. Rev. Lett. **61**, 826 (1988).
- [3] H. Wu, M. J. Holland, and C. J. Foot, J. Phys. **B 28**, 5025 (1995).
- [4] M. S. Zubairy, A. B. Matsko, and M. O. Scully, Phys. Rev. **A 65**, 043804 (2002).
- [5] L. Liu, J. Bellanca, and H. Metcalf, Phys. Rev. **A 65**, 043401 (2002).
- [6] K. J. Boller and A. Imamoglu, Phys. Rev. Lett. **66**, 2539 (1991).
- [7] S. E. Harris, Phys. Today **50**, 36 (1997).
- [8] M. O. Scully and M. S. Zubiary, *Quantum Optics* (Cambridge University Press, Cambridge, 1997).
- [9] M. D. Lukin and A. Imamoglu, Nature (London) **413**, 273 (2001).
- [10] J. I. Cirac, P. Zoller, H. J. Kimble, and H. Mabuchi, Phys. Rev. Lett. **78**, 3221 (1997).
- [11] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
- [12] D. Bouwmeester, A. Ekert and A. Zeilinger, eds., *The Physics of Quantum Information* (Springer Verlag, NY, 2000).
- [13] A. Mair, J. Hager, D. F. Phillips, R. L. Wadsworth, and M. D. Lukin, Phys. Rev. **A 65**, 031802 (2002).
- [14] A. Raichenbeutel, P. Bertet, S. Osnaghi, G. Nogues, M. Brune, J. M. Raimond, and S. Haroche, Phys. Rev. **A 64**, 050301 (2001).
- [15] Ian Walmsey and Peter Knight, Optics and Photonics News, p.43 (November 2000).
- [16] D. C. Roberts, T. Gasenzer, and K. Burnett, Phys. Rev. **A 66**, 023801 (2002).
- [17] W. S. Warren, H. Rabitz and M. Dahleh, Science **259**, 1581 (1993).
- [18] P. Král, Z. Amitay, and M. Shapiro, Phys.Rev.Lett. **89**, 063002 (2002).
- [19] B. W. Shore, *Theory of Coherent Atomic Excitation* (John Wiley, NY, 1990).
- [20] P. Milonni and J. H. Eberly, *Lasers* (John Wiley, NY, 1985).
- [21] P. Meystre, *Atom Optics* (Springer Verlag, NY, 2001).
- [22] L. Allen and J. H. Eberly, *Optical Resonance and Two-level Atoms* (Dover Publ., New York, 1987).
- [23] S. Guerin, F. Monti, J. M. Dupont and H. R. Jauslin, J. Phys. **A 30**, 7193 (1997).
- [24] Z. Bialynicka-Birula, I. Bialynicka-Birula, J. H. Eberly, and B. W. Shore, Phys. Rev. **A 5**, 2048 (1977).

- [25] Kh. Kh. Shakov and J. H. McGuire, Phys. Rev. **A 67**, 033405 (2003).
- [26] J. H. McGuire, Kh. Kh. Shakov, and Kh. Yu. Rakhimov, J. Phys. **B**, in press.
- [27] M. T. Frey, F. B. Dunning, C. O. Reinhold, S. Yoshida, and J. Burgdoerfer, Phys. Rev. **A 59**, 1434 (1999).
- [28] E. Persson, S. Yoshida, X.-M. Tang, C. Reinhold, and J. Burgdoerfer, Phys. Rev. **A 66**, 043407 (2002).
- [29] C. L. Stokley, F. B. Dunning, C. O. Reinhold, and A. K. Pattanayak, Phys. Rev. **A 65**, 021405(R) (2002).
- [30] W. Magnus, Commun. Pure and Applied Math **7**, 971 (1954).
- [31] J. H. McGuire and P. R. Simony, Phys. Rev. **A 22**, 2270 (1980).
- [32] J. H. McGuire, D. J. Land, J. G. Brennan, and G. Basbas, Phys. Rev. **A 19**, 2180 (1979).
- [33] T. F. Gallagher, *Rydberg Atoms* (Cambridge Univ. Press, Cambridge, 1994).
- [34] R. G. Hulet and D. Kleppner, Phys. Rev. Lett. **51**, 1436 (1983).
- [35] J. Lambert, M. W. Noel, and T. F. Gallagher, Phys. Rev. **A 66**, 053413 (2002).